

Headline AMP®

Fungicide

BASF

We create chemistry

FEATURED STORY
Read All About It


A Legacy of Success

When you need high-quality disease protection at a value price, there is no better option than Headline AMP® fungicide. For more than 10 years it has consistently proven its ability to fight disease, provide Plant Health benefits, and help increase yields. With a product as good as this, it's best to let the the results tell the story.

Benefits of Headline AMP Fungicide


- Highest yielding fungicide in its class
- Strong disease prevention and control
- Reduced plant stress and greater photosynthesis
- Plant Health benefits in all environmental conditions

Headline AMP Fungicide Outperforms the Competition



Summary of BASF or partially/fully sponsored University or Consultant small-plot replicated trials, 2014-2020. Headline AMP fungicide 10 fl oz/A, Aproach Prima fungicide 6.8 fl oz/A, Stratego YLD fungicide 4 fl oz/A, Quilt Xcel fungicide 10.5 fl oz/A.

Gray Leaf Spot Control in Field Corn




BASF Replicated Trial, Andrews, NC, 2013, R1 Application; Photo 28 DAT. Headline AMP fungicide applied at 10 fl oz/A, Quilt Xcel at 10.5 fl oz/A, Stratego YLD at 4 fl oz/A, Aproach at 6 fl oz/A.


Headline AMP[®] Fungicide Improved Canopy Health



Best-in-Class Northern Corn Leaf Blight Control



BASF sponsored replicated trial, sweet corn, U of Florida, 2013, 3 applications (late whorl stage, silk and silk + 7d). Harvested May 25.



BASF small-plot corn trial, Iowa, 2016 (RCB design, 4 replicates). Headline AMP fungicide applied at VT at 10 fl oz/A. Canopy photos taken 50 DAT. Lower canopy photos shown.

Apply Headline AMP Fungicide and Pay Nothing Until After Harvest

With the 2021 Grower Finance Program you can treat your fields using BASF's premium corn fungicide and pay absolutely nothing until December 2021¹. As part of the deal you can include select tank mix partners and your application costs.

Talk to your BASF Authorized Retailer to get started. See full program offer for details.



¹ Subject to credit approval, terms and conditions. Credit must be approved and transaction submitted by listed program dates in order to qualify for incentive rate. Offer valid on qualifying purchases made between October 1, 2020, to September 30, 2021.

Headline AMP®

Fungicide

Corn Fungicide Tech Sheet

Best Use Recommendations

| Crop* | Recommended Timing |
|---------------------|---|
| Corn (all types) | VT-R2: For optimal Plant Health benefits, including disease control and environmental stress mitigation |

| Crop | Diseases Controlled** | Application Rate |
|------|--|--------------------|
| Corn | <ul style="list-style-type: none">■ Anthracnose (<i>Colletotrichum acutatum</i>, <i>C. gloeosporioides</i>)■ Eyespot (<i>Aureobasidium zeae</i>)■ Gray leaf spot (<i>Cercospora zeae-maydis</i>)■ Northern corn leaf blight (<i>Exserohilum turcicum</i>)■ Rust, common (<i>Puccinia sorghi</i>)■ Rust, southern (<i>Puccinia polyspora</i>)■ Southern corn leaf blight (<i>Bipolaris maydis</i>)■ Tar spot (<i>Phyllachora maydis</i>) | 10-14.4 fl oz/A |

* See label for complete listing of labeled crops

** See label for complete listing of diseases controlled

Active Ingredients:

Metconazole – Group 3
Pyraclostrobin – Group 11

Formulations:

Suspension concentrate (SC)

Product Packaging:

2 x 2.5 gallons
Bulk options

Storage:

Store above 32°F

Bulk Density:

8.846 lbs/gal



■ BASF

We create chemistry

Application Tips

Rainfast: When dry

REI: 12 hours

PHI: 20 days for grain and stover, 7 days for sweet corn, silage and forage

Adjuvant: Can be used to improve coverage†

Nozzle, droplet size: Flat fan, fine to medium

† See label for adjuvant use limitation on corn

Water Volume

Aerial: Minimum 2 gpa spray volume

Ground: Minimum 10 gpa spray volume

Chemigation: **DO NOT** apply more than 1/2 inch per acre

DO NOT apply in spray solutions less than 50% water by volume

Adjuvant Use Limitation on Corn

Adjuvant crop damage can occur when an adjuvant is used after the V8 stage and before the VT stage (the VT stage is defined as when the tassel's last branch is completely visible outside the whorl). If an adjuvant is used after the V8 stage and before the VT stage, the grower and the user are responsible for contacting the adjuvant source (adjuvant distributor, retailer, or manufacturer) for advice and confirmation that the adjuvant has been tested and proved to be safe for application from the V8 to VT corn stages. Refer to the adjuvant label for specific use directions and restrictions. Always follow the most restrictive label.

Tank Mixes and Additives

Headline AMP® fungicide can be tank mixed with other fungicides, insecticides, herbicides, liquid fertilizers, biological control products, adjuvants, and additives. Always follow the most restrictive label use directions. Consult a BASF representative or local agricultural authorities for more information on use of additives or adjuvants with this product.

Mixing Order

1. **Water:** Begin by agitating a thoroughly clean sprayer tank $\frac{3}{4}$ full of clean water
2. **Agitation:** Maintain constant agitation throughout mixing and application
3. **Products in PVA bags**
4. **Water-dispersible products** (products such as dry flowables, wettable powders, suspension concentrates such as **Headline AMP fungicide**, or suspo-emulsions)
5. **Water-soluble products**
6. **Emulsifiable concentrates** (or oil concentrates when applicable)
7. **Water-soluble additives** (such as ammonium sulfate [AMS] or urea ammonium nitrate [UAN] when applicable)
8. **Nutritionals**
9. **Remaining quantity of water**



To learn more about crop protection products from BASF, visit www.agproducts.basf.us

 **BASF**

We create chemistry

Headline AMP®
Fungicide