

Soybean Fungicide Tech Sheet

Best Use Recommendations

Crop*

Recommended Timing

Soybean

R2-R4:

For optimal Plant Health benefits, including disease control and environmental stress mitigation

Crop	Diseases Controlled†	Application Rate
Soybean	<ul style="list-style-type: none"> ■ Alternaria leaf spot (<i>Alternaria</i> spp.) ■ Anthracnose (<i>Colletotrichum truncatum</i>) ■ Asian soybean rust (<i>Phakopsora pachyrhizi</i>) ■ Brown spot (<i>Septoria glycines</i>) ■ Cercospora blight (<i>Cercospora kikuchii</i>) ■ Frogeye leaf spot (<i>Cercospora sojina</i>)‡ ■ Pod and stem blight (<i>Diaporthe phaseolorum</i>) ■ Rhizoctonia aerial blight (<i>Rhizoctonia solani</i>) ■ Target spot (<i>Corynespora cassicola</i>) ■ Sclerotinia blight (white mold) (<i>Sclerotinia sclerotiorum</i>)§ 	4 fl oz/A
	<ul style="list-style-type: none"> ■ Southern blight (<i>Sclerotium rolfsii</i>)§ 	8 fl oz/A

* See label for complete listing of labeled crops

† See label for complete listing of diseases controlled

‡ Addition of a triazole fungicide may be required for control of QoI-resistant frogeye leaf spot

§ Suppression only

Active Ingredients:

Fluxapyroxad – Group 7
Pyraclostrobin – Group 11

Formulation:

Suspension concentrate (SC)

Product Packaging:

2 x 1 gallon
2 x 2.5 gallons
110 gallon mini bulk

Storage:

Store above 32°F

Bulk Density:

9.6807 lbs/gal



■ BASF

We create chemistry

Application Tips

Rainfast: When dry

REI: 12 hours

PHI: 21 days for soybean, except; 14 days for soybean forage

Adjuvant: Can be used to improve coverage

Nozzle, droplet size: Flat fan, fine to medium

Water Volume

Aerial: Minimum 2 gpa spray volume

Ground: Minimum 10 gpa spray volume

Chemigation: **DO NOT** apply more than 1/2 inch per acre

DO NOT apply in spray solutions less than 50% water by volume



Tank Mixes and Additives

Priaxor® fungicide can be tank mixed with other fungicides, insecticides, herbicides, liquid fertilizers, biological control products, adjuvants, and additives. Always follow the most restrictive label use directions. Consult a BASF representative or local agricultural authorities for more information on use of additives or adjuvants with this product.

Mixing Order

1. **Water:** Begin by agitating a thoroughly clean sprayer tank $\frac{3}{4}$ full of clean water
2. **Agitation:** Maintain constant agitation throughout mixing and application
3. **Products in PVA bags**
4. **Water-dispersible products** (products such as dry flowables, wettable powders, suspension concentrates such as **Priaxor fungicide**, or suspo-emulsions)
5. **Water-soluble products**
6. **Emulsifiable concentrates** (or oil concentrates when applicable)
7. **Water-soluble additives** (such as ammonium sulfate [AMS] or urea ammonium nitrate [UAN] when applicable)
8. **Nutritionals**
9. **Remaining quantity of water**



To learn more about crop protection products from BASF, visit www.agproducts.basf.us

 **BASF**

We create chemistry

Priaxor®

Xemium® Brand Fungicide